

National Registry of Emergency Medical Technicians Advanced Level Psychomotor Examination

SUPRAGLOTTIC AIRWAY DEVICE

Candidate:	Examiner:		
	Signature:		
Device:			
	ally ventilate with BVM attached to reservoir and oxygen, full credit must be a ation is delivered within 30 seconds.	warded for step	os denoted by
Actual Time Started:	_	Possible Points	Points Awarded
Takes or verbalizes body substance isolation precautions		1	
Opens the airway manually		1	
Elevates tongue, inserts simple adjunct [oropharyngeal or nasopharyngeal airway]		1	
NOTE: Examiner now informs of	candidate no gag reflex is present and patient accepts adjunct		
**Ventilates patient immediately with bag-valve-mask device unattached to oxygen		1	
**Ventilates patient with room air		1	
NOTE: Examiner now informs of the patient's blood oxygen satu	candidate that ventilation is being performed without difficulty and that μ	oulse oximetry	indicates
Attaches oxygen reservoir to bag [12 – 15 L/minute]	-valve-mask device and connects to high-flow oxygen regulator	1	
Ventilates patient at a rate of 10 -	- 12/minute with appropriate volumes	1	
	miner auscultates and reports breath sounds are present and equal of a supraglottic airway. The examiner must now take over ventilation.	bilaterally an	nd medical
Directs assistant to pre-oxygenate	e patient	1	
Checks/prepares supraglottic airv	way device	1	
Lubricates distal tip of the device	[may be verbalized]	1	
NOTE: Examiner to remove OP	A and move out of the way when candidate is prepared to insert device.		
Positions head properly		1	
Performs a tongue-jaw lift		1	
Inserts device to proper depth		1	
Secures device in patient [inflates cuffs with proper volumes and immediately removes syringe or secures strap]		1	
Ventilates patient and confirms problems bilaterally over lungs and over ep	roper ventilation [correct lumen and proper insertion depth] by auscultation igastrium	1	
Adjusts ventilation as necessary [ventilation is optimized]	ventilates through additional lumen or slightly withdraws tube until	1	
Verifies proper tube placement by colorimetric device	y secondary confirmation such as capnography, capnometry, EDD or	1	
NOTE: The examiner must now each ventilation?"	ask the candidate, "How would you know if you are delivering appropri	iate volumes v	vith
Secures device or confirms that the device remains properly secured		1	
Ventilates patient at proper rate and volume while observing capnography/capnometry and pulse oximeter		1	
Actual Time Ended:	TOTA	L 19	

Failure to initiate ventilations within 30 seconds after taking body substance isolation precautions or interrupts ventilations for greater than
seconds at any time
Failure to take or verbalize body substance isolation precautions
Failure to voice and ultimately provide high oxygen concentration [at least 85%]
Failure to ventilate the patient at a rate of 10 – 12/minute
Failure to provide adequate volumes per breath [maximum 2 errors/minute permissible]
Failure to pre-oxygenate patient prior to insertion of the supraglottic airway device
Failure to insert the supraglottic airway device at a proper depth or location within 3 attempts
Failure to inflate cuffs properly and immediately remove the syringe
Failure to secure the strap (if present) prior to cuff inflation
Failure to confirm that patient is being ventilated properly (correct lumen and proper insertion depth) by auscultation bilaterally over lungs a
over epigastrium
Insertion or use of any adjunct in a manner dangerous to the patient
Failure to manage the patient as a competent EMT
Exhibits unacceptable affect with patient or other personnel
Uses or orders a dangerous or inappropriate intervention

You must factually document your rationale for checking any of the above critical items on the reverse side of this form.

© 2011 National Registry of Emergency Medical Technicians, Inc., Columbus, OH
All materials subject to this copyright may be photocopied for the non-commercial purpose of educational or scientific advancement.

Critical Criteria

p304/06-12